

# Package: extRC (via r-universe)

August 21, 2024

**Type** Package

**Title** Extended RC Models for Contingency Tables

**Version** 1.2

**Date** 2020-10-10

**Author** Francesco Bartolucci, Antonio Forcina

**Maintainer** Francesco Bartolucci <francesco.bartolucci@unipg.it>

**Description** Maximum likelihood estimation of an extended class of row-column (RC) association models for two-dimensional contingency tables, which are formulated by a condition of reduced rank on a matrix of extended association parameters; see Forcina (2019) <[arXiv:1910.13848](#)>. These parameters are defined by choosing the logit type for the row and column variables among four different options and a transformation derived from suitable divergence measures.

**License** GPL (>= 2)

**Imports** MASS

**NeedsCompilation** no

**Date/Publication** 2020-10-10 21:00:02 UTC

**Repository** <https://francescobartolucci.r-universe.dev>

**RemoteUrl** <https://github.com/cran/extRC>

**RemoteRef** HEAD

**RemoteSha** 716b900b34bac7de59675cba979c84f78f951820

## Contents

extRC-package	2
cuby	3
Deta	4
dfm	5
Drank	5
extRC	6

Hmat	7
MainRC	8
MatIn	9
mobility	9
plot	10
PraD	11
print	11
summary	12
tril	12
<b>Index</b>	<b>14</b>

---

extRC-package	<i>Overview of the Package extRC</i>
---------------	--------------------------------------

---

## Description

Estimation of extended RC models, which are formulate by constraining different types of association parameters to have a reduced rank.

## Details

The package contains functions for maximum likelihood (ML) estimation of an extended class of row-column (RC) association models for two-dimensional contingency tables, as described in Forcina (2019). These models are formulated by a condition of reduced rank on a matrix of extended association parameters, which are defined by choosing the logit type for the row and column variables among four different options and a transformation derived from Cressie and Read (1984). Among the available alternatives, it is possible to use log-odds ratio based on different types of aggregation of the joint probabilities. The class of models generalizes that proposed in Kateri and Papaioannou (1994), Bartolucci and Forcina (2002), and Esendiller (2017), and includes the original RC association models of Goodman (1979) and the correspondence analysis model, as formulated in Goodman (1981) and Gilula et al. (1988). Maximum likelihood estimation is based on an algorithm that is an adaptation of the Aitchison and Silvey (1958) algorithm for constrained ML estimation and is related to the algorithm described in Evans and Forcina (2013) for fitting constrained marginal models.

The main function in the package is `extRC` that provides an output that may be shown by usual R commands `print`, `summary`, and `plot`.

## Author(s)

Francesco Bartolucci, Antonio Forcina

Maintainer: Francesco Bartolucci <francesco.bartolucci@unipg.it>

## References

- Aitchison, J. and Silvey (1958). Maximum-likelihood estimation of parameters subject to restraints. *The Annals of Mathematical Statistics*, **29**, 813-828.
- Bartolucci, F. and Forcina, A. (2002). Extended RC association models allowing for order restrictions and marginal modeling. *Journal of the American Statistical Association*, **97**, 1192-1199.
- Cressie, N. and Read, T.R., 1984. Multinomial goodness-of-fit tests. *Journal of the Royal Statistical Society: Series B*, **46**, 440-464.
- Espendiller, M., 2017. *Association in contingency tables*. Ph.D. thesis.
- Evans, R.J. and Forcina, A. (2013). Two algorithms for fitting constrained marginal models. *Computational statistics & Data analysis*, **66**, 1-7.
- Forcina (2019), An extended class of RC association models: definition and estimation, arXiv:1910.13848.
- Gilula, Z., Krieger, A.M., and Ritov, Y., 1988. Ordinal association in contingency tables: Some interpretive aspects. *Journal of the American Statistical Association*, **83**, 540-545.
- Goodman, L.A. (1979). Simple models for the analysis of association in cross-classifications having ordered categories. *Journal of the American Statistical Association*, **74**, 537-552.
- Goodman, L.A., 1981. Association models and canonical correlation in the analysis of cross-classifications having ordered categories. *Journal of the American Statistical Association*, **76**, 320-334.
- Kateri, M. and Papaioannou, T. (1994). *f-divergence Association Models*. University of Ioannina.

## Examples

```
# load data
data(mobility)

# fit model for a single la
out = extRC(mobility,mod=c("1","1"),k=1,la=0.6)
summary(out)
```

---

cuby

*Step length*

---

## Description

Internal function that computes step length of the estimation algorithm in [extRC](#) by fitting a cubic polynomial.

## Usage

```
cuby(g)
```

## Arguments

**g** vector of likelihood values at different step lengths

**Value**

comp1            optimal length

**Author(s)**

Francesco Bartolucci, Antonio Forcina

---

Deta

*Computation of marginal parameters*

---

**Description**

Given a vector of canonical parameters coding distribution for an  $I \times J$  contingency table and the RC model specification in list Model, it computes vector of marginal and joint parameters and matrix of its derivatives with respect the canonical parameters.

**Usage**

Deta(the, Model, der = FALSE)

**Arguments**

the            vector of canonical parameters  
 Model        list specifying all model components  
 der           to require derivative computation (optional)

**Value**

eta            vector of marginal parameters  
 Der           derivative matrix with respect to canonical parameters

**Author(s)**

Francesco Bartolucci, Antonio Forcina

---

dfm	<i>First difference matrix</i>
-----	--------------------------------

---

**Description**

It creates a matrix of first differences of order  $k$ .

**Usage**

```
dfm(k)
```

**Arguments**

k	size of the matrix
---	--------------------

**Value**

D	first difference matrix
---	-------------------------

**Author(s)**

Francesco Bartolucci, Antonio Forcina

**Examples**

```
D = dfm(5)
x = runif(5)
(D*%x)
```

---

Drank	<i>Check matrix rank</i>
-------	--------------------------

---

**Description**

Given the row vectorized matrix, it computes the vector of discrepancies with respect to a certain rank and its derivative.

**Usage**

```
Drank(ga, lev, k, der = FALSE)
```

**Arguments**

ga	row vectorized matrix of interaction
lev	vector of the number of row and column categories in the original table (the numbers of rows and columns of the input matrix must be increased by 1)
k	matrix rank
der	to require derivative

**Value**

fr                    vector of discrepancies with respect to the rank  
 Dfr                   derivative of fr

**References**

Bartolucci, F. and Forcina, A. (2002). Extended RC association models allowing for order restrictions and marginal modeling. *Journal of the American Statistical Association*, **97**, 1192-1199.

**Examples**

```
A = matrix(rnorm(12),4) # matrix the rank of which must be checked
a = as.vector(t(A))
out = Drank(a,c(5,4),1,der=TRUE)
(out$fr)
(out$Dfr)
```

---

 extRC

---

*Extended RC model*


---

**Description**

Main function that fits extended RC models based on different types of aggregation (continuation, local, global) and different divergence functions defined by a suitable value of lambda.

**Usage**

```
extRC(N, mod, k, la, marg.cons = c("free", "equal", "shift"))
```

**Arguments**

N                    observed contingency table  
 mod                vector indicating the types of aggregation for row and column variables ("c" for continuation, "l" for local, "g" for global)  
 k                    rank required for the matrix of interaction parameters  
 la                  value of lambda parameter  
 marg.cons        type of constraint on the marginal distributions

**Value**

la                   vector of lambda values (when a vector is in input)  
 dev                deviance of the fitted model (when only one lambda value is in input) or vector of deviances (when a vector of lambda values is in input)  
 df                  degrees of freedom (when only one lambda value is in input)  
 it                  number of iterations (when only one lambda value is in input)

dis	final discrepancy (when only one lambda value is in input)
pj	vector of joint probabilities under the fitted model (when only one lambda value is in input)
eta	full vector of marginal parameters (when only one lambda value is in input)
etaX	vector of row marginal parameters (when only one lambda value is in input)
etaY	vector of column marginal parameters (when only one lambda value is in input)
Eta	matrix of association parameters (when only one lambda value is in input)
la	vector of lambda values (when more lambda values are in input)
dev	vector of deviance values (when more lambda values are in input)

**Author(s)**

Francesco Bartolucci, Antonio Forcina

**Examples**

```
# load data
data(mobility)

# for a single value of lambda, fit model with constraints of rank 1 on
# local-local logits and without constraints on the marginal distributions
out = extRC(mobility,mod=c("l","l"),k=1,la=0.6)
summary(out)

# for a single value of lambda, fit model with constraints of rank 1 on
# local-local logits and under constrain of equal marginal distributions
out = extRC(mobility,mod=c("l","l"),k=1,la=0.6,marg.cons="equal")
summary(out)

# for a single value of lambda, fit model with constraints of rank 2 on
# global-global logits and under constraint that marginal distributions
# are equal up to a constant shift
out = extRC(mobility,mod=c("g","g"),k=2,la=0.6,marg.cons="shift")
summary(out)

# fit model for a vector of lambdas
la = seq(-1.8,0.6,length.out=10)
out1 = extRC(mobility,mod=c("l","l"),k=1,la=la)
plot(out1)
```

---

Hmat

*Matrix algebra transformation*


---

**Description**

Internal function that performs a matrix algebra transformation that is used for estimation in [extRC](#).

**Usage**`Hmat(G)`**Arguments**`G` input matrix**Value**`H` transformed matrix**Author(s)**

Francesco Bartolucci, Antonio Forcina

---

`MainRC`*Estimation of extended RC models*

---

**Description**

Internal function that implements the Aitchinson-Silvey algorithm to estimate extended RC models.

**Usage**`MainRC(y, Model, the0 = NULL, output = FALSE)`**Arguments**`y` row vectorized vector of frequencies of the contingency table`Model` list of model components`the0` initial vector of canonical parameters (optional)`output` to require full output (optional)**Value**`dev` final deviance`df` degrees of freedom`pj` vector of joint probabilities`it` number of iterations`dis` final discrepancy**Author(s)**

Francesco Bartolucci, Antonio Forcina

---

MatIn	<i>Aggregation matrices</i>
-------	-----------------------------

---

**Description**

Computation of aggregation matrices for generalized interactions that are used in `codeextRC` to estimate extended RC models.

**Usage**

```
MatIn(lev, mod)
```

**Arguments**

lev	vector number of rows and columns
mod	type of logit for each dimension

**Value**

R0	aggregation matrix for the row margin upper level
R1	aggregation matrix for the row margin lower level
C0	aggregation matrix for the column margin upper level
C1	aggregation matrix for the column margin lower level
J00	aggregation matrix for the left upper quadrant
J01	aggregation matrix for the right upper quadrant
J10	aggregation matrix for the left lower quadrant
J11	aggregation matrix for the right lower quadrant

**Author(s)**

Francesco Bartolucci, Antonio Forcina

---

mobility	<i>Social mobility data</i>
----------	-----------------------------

---

**Description**

Social mobility table of 3,500 British individuals, who are cross-classified according to their occupational status and the occupation status of their fathers.

**Usage**

```
data("mobility")
```

**Format**

The format is: num [1:5, 1:5] 50 28 11 14 3 45 174 78 150 42 ... - attr(\*, "dimnames")=List of 2 ..\$ : chr [1:5] "F1" "F2" "F3" "F4" ... ..\$ : chr [1:5] "S1" "S2" "S3" "S4" ...

**References**

Mosteller, F. (1968). Association and estimation in contingency tables. *Journal of the American Statistical Association*, **63**, 1-28.

Bartolucci, F. and Forcina, A. (2002). Extended RC association models allowing for order restrictions and marginal modeling. *Journal of the American Statistical Association*, **97**, 1192-1199.

---

 plot

*Plot the output*


---

**Description**

It plots the output of `codeextRC` function for a vector of lambda values.

**Usage**

```
## S3 method for class 'extRC'
plot(x, ...)
```

**Arguments**

x                    output from `extRC`  
 ...                  further arguments passed to or from other methods

**Value**

None

**Author(s)**

Francesco Bartolucci, Antonio Forcina

---

PraD	<i>Discrepancy with respect to equality constraints</i>
------	---

---

**Description**

Internal function that, given a vector of canonical parameters for an  $I \times J$  table (vectorized by row) and the RC model specification in list Model, computes vector of discrepancies and matrix of its derivatives with respect to the canonical parameters.

**Usage**

```
PraD(the, Model, der = FALSE)
```

**Arguments**

the	vector of canonical parameters
Model	list of model components
der	to require the derivative (optional)

**Value**

eta	vector of parameters (logits, interactions)
hdis	vector of discrepancies
Hdis	matrix of derivatives of discrepancies with respect to the canonical parameter (optional)

**Author(s)**

Francesco Bartolucci, Antonio Forcina

---

print	<i>Print the output.</i>
-------	--------------------------

---

**Description**

Given the output of `codeextRC` function, it is written in a readable form.

**Usage**

```
## S3 method for class 'extRC'
print(x, ...)
```

**Arguments**

x	output from <code>extRC</code>
...	further arguments passed to or from other methods

**Value**

None

**Author(s)**

Francesco Bartolucci, Antonio Forcina

---

summary	<i>Summary of extRC fits</i>
---------	------------------------------

---

**Description**Summary method for the output of `codeextRC` function.**Usage**

```
## S3 method for class 'extRC'  
summary(object, ...)
```

**Arguments**

object	output from <code>extRC</code>
...	further arguments passed to or from other methods

**Value**

None

**Author(s)**

Francesco Bartolucci, Antonio Forcina

---

tril	<i>Lower triangular matrix</i>
------	--------------------------------

---

**Description**

Given a square matrix, it provides the lower triangular part, including the main diagonal.

**Usage**

```
tril(M)
```

**Arguments**

M	square matrix
---	---------------

**Value**

N                    transformed matrix

**Author(s)**

Francesco Bartolucci, Antonio Forcina

**Examples**

```
M = matrix(1:9,3)
N = tril(M)
```

# Index

- \* **algebra**
  - Drank, [5](#)
  - Hmat, [7](#)
- \* **array**
  - tril, [12](#)
- \* **datasets**
  - mobility, [9](#)
- \* **multivariate**
  - extRC-package, [2](#)

cuby, [3](#)

Deta, [4](#)  
dfm, [5](#)  
Drank, [5](#)

extRC, [2](#), [3](#), [6](#), [7](#), [9–12](#)  
extRC-package, [2](#)

Hmat, [7](#)

MainRC, [8](#)  
MatIn, [9](#)  
mobility, [9](#)

plot, [2](#), [10](#)  
PraD, [11](#)  
print, [2](#), [11](#)

summary, [2](#), [12](#)

tril, [12](#)